

Sensitivity of upland arthropod diversity to livestock grazing, vegetation structure and landform

Peter Dennis

Macaulay Institute, Craigiebuckler, Aberdeen, AB15 8QH, Scotland, UK. e-mail: p.dennis@macaulay.ac.uk

Received 13 January 2003, accepted 30 April 2003.

Abstract

Livestock production extends to most non-forested, marginal, upland habitats of Britain. Of these, indigenous grasslands are stocked predominantly by sheep, stocking densities having increased in Scotland by 25% between 1975 and 1990. Conversely, the national herd of cattle in Scotland declined by 22% over the same period. The effects of grazing management on arthropod distribution and abundance is reviewed, with particular emphasis on the results of grazing experiments that have investigated the effects on arthropods of varied livestock species and stocking rates. Arthropods contribute the most species of any taxa in the uplands and are critical in upland food chains. The direction and magnitude of the response of different arthropod taxa to grazing management reflects their trophic level, life history, size and mobility, e.g., 30% of ground and rove beetle species are more sensitive to landform than grazing management. For the arthropod taxa that are sensitive to grazing management, the effects are generally indirect, via changes in the heterogeneity of botanical composition and vegetation structure. A mosaic of contrasting botanical composition and structural heterogeneity is essential to conserve and enhance arthropod and broader wildlife diversity in the uplands. However, the landscape-scale study of mammalian herbivore-vegetation-arthropod interactions is required both to quantify the relative importance of land use (grazing management) and landform (landscape physiognomy) across the uplands and to determine the optimal grain-size of the habitat mosaic to sustain biodiversity.

Key words: Livestock production, sheep grazing, cattle grazing, indigenous grasslands, upland, biodiversity, insect diversity and spider diversity.

Introduction

Substantial increases in Ovis aries (sheep) numbers in upland areas of Britain from the 1970s-1990s (2 million or 25%) have placed significant pressure on indigenous grasslands in the uplands ¹. There has been a 600 000 or 22% decline in Bos taurus (cattle) across all farmland over the same period, indicating a trend from mixed grazers to sheep only. Grazing beyond sustainable intensities by sheep, Cervus elaphus (red deer) and Oryctolagus cuniculus (rabbits) has also increased grassland at the expense of heathland vegetation dominated by Calluna vulgaris (heather) ^{2,3,4,5,6}. Hence, many uplands are currently grass- rather than heather-dominated ⁷ although the arthropod fauna of indigenous grasslands can also be diverse compared with alternative biotopes ^{8,9}. Indigenous upland grasslands also have a greater importance today because the species shared with lowland grasslands have been diminished by intensification of management in the lowlands¹⁰ Upland grasslands vary in their botanical composition although there is a predominance of acid grasslands composed of Agrostis and Festuca species with dominance by Nardus stricta in drier areas and Molinia caerulea, Eriophorum or Scirpus species in wetter areas ¹¹. Research on grassland arthropods has largely focused on lowland, calcareous grasslands because of the higher botanical diversity compared with upland, acid grasslands ¹² and the distinct arthropod species rather than diversity per se 12,9. However, there are substantial areas of karst and small calcareous flushes in the uplands and these have an important arthropod fauna associated with them ^{13,14}. The extensification of pasture management is considered a means of integrating biodiversity conservation with agricultural production objectives for the broader upland grassland types and for achieving a restoration of heather moorlands 15.

Arthropod diversity - so what?

Typically, arthropods contribute well over half the metazoan species in any habitat or ecosystem ¹⁶. Beyond the numbers argument,

arthropods occupy several trophic levels and fulfil vital functions within ecosystems that can be summarized under particular socioecological headings ^{17,18}. Certain species have utility value as natural enemies of potential pest herbivores in grasslands and heathlands. For example, the ground beetles, Carabus spp. and Cychrus caraboides (Coleoptera: Carabidae) are mollusc predators, the smaller but closely related Notiophilus aquaticus specializes on springtails (Collembola) that includes the pest, Sminthurus viridis ¹⁹. Many herbivorous insects are not pests but they do consume plant material and return nutrients to the soil, hence reducing litter accumulation. They also provide food for higher trophic levels (predatory insects, insectivorous mammals and birds). Some Diptera (frit flies and craneflies), Coleoptera (leaf beetles, weevils and wireworms) and springtails (Sminthurus viridis) can be occasional pests in intensively managed grasslands although this is unknown in indigenous grasslands.

There has been more general scientific interest (ecologisticscientific value) in what arthropod species do in ecosystems ¹⁸, particularly inspired by evidence of declines in abundance of many species and questions of what the consequences might be for ecosystem function. The productivity and maintenance of fertility in soils is one critical area, dung beetles prevent pasture fouling and enhance the cycling of nutrients, whilst bumblebees ensure the successful pollination of many flowering species. The ecologisticscientific value extends to the role for arthropods as contributors to the diet of other species of conservation value. Leatherjackets, larvae of craneflies (Diptera: Tipulidae), represent the largest invertebrate biomass in the uplands and significantly contribute to the diet of the rare upland Dotterel, Eudromias morinellus (Charadriidae) ²⁰ and the economically important Red grouse, Lagopus lagopus ²¹. Large beetles, such as Carabus spp. (Col.: Carabidae) and Dor beetles (Col.: Scarabaeidae) make a contribution to the diet of birds of prey such as the Merlin, Falco columbarius (Falconidae) 22. Finally, the naturalistic, aesthetic value of the more charismatic species of insects is self-evident

but highly selective. The Scotch Argus butterfly, *Erebia aethiops* (Lepidoptera: Satyridae), Emperor moth, *Saturnia pavonia* (Lep.: Saturnidae), Burnet moths, *Zygaena* spp. (Lep.: Zygaenidae), and dragonflies and darters, *Aeshna juncea* and *Sympetrum danae* (Odonata) are typical examples (UK Biodiversity Action Plan¹⁶).

The importance of Spatial Scale

It is important to take account of the ways in which spatial scale impinges on large herbivore-arthropod interactions. Certainly, large herbivores interact with vegetation at three identifiable scales: the feeding station, vegetation patch and landscape ²³. The relative effects of large herbivores on arthropods varies according to the scale of observation because:

• large herbivores do not uniformly forage everywhere ^{4,24}.

• plant communities, and their availability and response to grazing ^{25,26}, are not constant due to variability in landform, hydrology and climate¹¹.

Aspect, gullying and wet pockets alter the dominance of vegetation at constant altitude around hills and, in addition, altitudinal gradients in upland terrain are related to shifts in vegetation communities from dry grasslands, to heather and weathered peatlands or montane vegetation on skeletal soil on the high plateaux ¹¹, all of which create different upland arthropod assemblages ^{27,28}. This paper reviews the findings from several experiments across Scotland that investigated arthropod responses to varied grazing management ^{29,30,31,32}. The results of the grazing experiments are placed in the context of other grazing studies of upland and indigenous habitats ^{33,34, 35,36, 37,38}.

Grazing Management on Upland Plant Communities

The research refers to several experiments that were conducted between 1993 and 1998 which had in common grazing by sheep, in one case cattle also, and all grazed to achieve a selection of target average sward heights, representing a spectrum of grazing intensities designed to match the productivity of each site. The site habitats defined according to Ratcliffe⁸ and given National Vegetation Classification (NVC) classes of Rodwell¹¹ included:

- Agrostis-Festuca equivalent to NVC U4e grazed by sheep to maintain 3, 4.5 and 6 cm sward heights, plus an ungrazed control, Cleish Hills, Fife.
- *Festuca-Agrostis* equivalent to NVC U4a grazed by sheep to maintain 3, 4.5 and 6 cm sward heights, plus an ungrazed control, Tyndrum, Highlands.
- *Nardus stricta* equivalent to NVC U5a grazed by sheep and cattle (June to August) to maintain 4.5 and 6.5 cm sward heights, plus an ungrazed control, Cheviot Hills, Scottish Borders.

Sampling arthropods and vegetation

Data collection of arthropods included the use of suction sampling for leafhoppers and web-spinning spiders on vegetation ³⁰, and pitfall trapping to sample surface-active beetles (Coleoptera) ^{29,31}, spiders (Araneae) and harvestmen (Opiliones) ³¹. Sward height profiles were measured around all arthropod-sampling points to provide data on spatial heterogeneity (patchiness) of the upland habitats ³⁰.

Distinguishing Direct/ Indirect Effects of Grazing Herbivores From Abiotic Influences on Arthropods

The interactions between grazing herbivores and vegetation and the magnitude of their influence on arthropods depends on geo-

Table 1. Turnover in dominant leafhopper and plant bug species of upland plant communities(• indicates presence).

	Plant community (NVC)	U4a <i>Festuca-</i>	U4e Agrostis-	U5a
_	Nardus	Agrostis	Festuca	stricta
	Leafhoppers - Homoptera Javasella discolor Dikraneura variata Diplocolenus abdominalis Aphrodes bifasciatus Plant bugs – Heteroptera Pachytomella parallela	•	•	•
Log abundance	3 2.5 2 1.5 1 0.5 0		■ 3.0 ■ 4.5 ■ 6.0 □ 12.0	cm cm cm) cm

Figure 1. Abundance of planthoppers (Homoptera: Auchenorrhyncha) in contrasting structural components of a patchy *Agrostis-Festuca* grassland (National Vegetation Classification U4e) grazed to different sward heights (after Dennis *et al*³⁰).

Sward

Hummock



Figure 2. Money spider responses to grazing management on a *Nardus stricta* grassland (S: sheep, C: cattle, UNG: ungrazed and numbers refer to target average inter-tussock sward height of treatment, cm).

graphical context. Complex landform is symptomatic of upland areas. Altitude, aspect and slope vary markedly to produce different patterns of insolation and soil wetness. For example, the topographic gradients of chalk grasslands revealed as much variance within-site as along a latitudinal gradient in England ³⁹. To place the effects of grazing herbivores on arthropods into context it is necessary to consider local biotic mechanisms and to interpret them with regard to broader environmental conditions.



Figure 3. Web numbers in microhabitats under different grazing regimes applied to a *Nardus stricta* grassland. Legend defined in Fig. 2 caption (afterDennis et al.³⁰).



Figure 4. Leafhopper and plant bug species responses to sheep grazing intensity on different plant communities.Key: triangles, *Festuca-Agrostis*, U4a; squares, *Agrostis- Festuca*, U4e and diamonds, *Nardus stricta*, U5a communities.

250 m

.

250 n



Figure 5. Size and location of clusters of high and low numbers of ground and rove beetles on *Nardus stricta* grassland under varied grazing management (after Dennis *et al.*,³²). Grazing treatments indicated on b, where S: sheep; C: cattle; UNG: ungrazed for 2-3 years; and, 4.5 and 6.5 represents average between-tussock sward height (cm). Upward pointed triangles represent aggregations of high value and upturned triangles aggregations of low numbers (where P_{G(z)} \propto size; Getis and Ord.⁶³).



Figure 6. The response of Coleoptera (Carabidae and Staphylinidae) species to grazing disturbance, topography and an interaction between these factors on upland *Nardus stricta* grassland. The 73 species of the assemblage were associated with each factor according to their intercorrelation in a community ordination and the scale and direction of response of indicator species using geostatistical analysis (Dennis *et al³⁰*). The satellite circles illustrate the direction of response of the species related to each factor.

Localised selection of preferred plant species

Small, insect herbivores, namely leafhoppers (Homoptera: Auchenorrhyncha) and other plant bugs (Heteroptera) select for particular species or groups of plants^{40, 41}. Particular species will therefore associate with upland communities where these host plants are abundant (Table 1) and direct competition between large herbivores and these insects will occur when grazing selects those host plants and reduces their abundance within a particular upland community ⁴².

Structural heterogeneity of vegetation and insect diversity

Grazing herbivores have an indirect effect on many arthropods by modifying the structural characteristics of upland habitats ³⁰. The structural appearance of vegetation can be dramatically altered by grazing regime, more so than plant species composition (Plate 1)^{43,44,45}. The development of distinct tussock or hummock and sward components has consequences for the distribution of small arthropods, both leafhoppers and spiders ³⁰. These structural components of grasslands contributed to arthropod species diversity. More individuals and species of leafhoppers were sampled in the taller, more complex components of grasslands, either tussock or hummocks (Fig. 1). Further, the impact of increased grazing intensity on these insects was, somewhat, buffered by these structures (Fig. 1).

If we consider predators requiring architectural diversity for the anchorage of webs, e.g., the money spider, *Silometopus elegans*, its general abundance increases where there is a lower grazing intensity but declines where grazing is absent for over two years (Fig. 2). By contrast, the common money spider, *Lepthyphantes mengii*, demonstrates a similar trend except that the highest catches were in the ungrazed *Nardus stricta* (Fig. 2). The differ-



Plate 1. Structural differences in *Nardus stricta* grassland after three years of contrasting grazing management, i. sheep grazed May to October to maintain a 6.5 cm inter-tussock sward, ii. sheep, May to October, with cattle, June to August, grazed to maintain a 4.5 cm inter-tussock sward.

ence is caused by the different locations of web building by these spiders. *Lepthyphantes mengii* constructs webs high in the leaves of tussocks or hummocks and this microhabitat is increased where grazing ceases for longer periods (Fig. 3). Taller swards were the most influential factor, typically being patchier, i.e., having greater variability in sward height ³⁰.

Effects of landform as opposed to solely grazing management The uplands represent complex terrain and it is important to disentangle direct/indirect effects of grazing management from variability imposed by landform. Slope and aspect create different patterns of insolation and soil wetness³⁹. Snow cover can be prolonged on north-facing slopes at higher altitude.

There was an interaction between grazing intensity and environmental conditions for leafhopper and plant bug species indicated by the consistent declining numbers with increased stocking density (Fig. 4). There were fewer potential species on the less productive vegetation at high altitude, on wet, cold slopes, e.g., *Festuca-Agrostis*. In addition, there was greater sensitivity of the plant bug species to grazing intensity on the sites of lower productivity (Fig. 4). Ground and rove beetles (Coleoptera: Carabidae, Staphylinidae), predatory species that roam around on the ground over tens of metres, were sampled at 120 points where pitfall traps were placed within the grazing experiment on *Nardus stricta* (U5a community). A geostatistical procedure was used to determine the size and location of clusters of high and low numbers of beetle species representative of the main trends of all the species²⁹. Four distinct patterns were identified that accounted for the main trends expressed by all species and the ecological interpretation identified the relative influences of landform and grazing management (Fig. 5). A large south-facing cluster that extended over several contrasting grazing treatments characterized Calathus melanocephalus (Col.: Carabidae; Fig. 5a), and indeed this species is typical of productive lowland pastures. There was also a large cluster across contrasting treatments for Philonthus decorus (Col.: Stapylinidae; Fig. 5c) and this showed the selection of more northern conditions typified by soil of greater wetness and organic matter content. The remaining two species both suggested responses to the patterns of grazing. Olophrum piceum (Col.: Staphylinidae) had clusters only within the ungrazed plots (Fig. 5b), and this probably relates to the increased litter and associated fungi that would develop in the absence of grazing. The clusters of high numbers of the large species, Carabus problematicus (Col.: Carabidae) associated with the sheep rather than sheep and cattle grazed plots (Fig. 5d), and this suggests that there are effects of cattle, possibly through soil compaction or direct treading disturbance, that are detrimental to this species. Carabus spp. depend on soil crevices as daytime refugia to avoid desiccation and predation, and the availability of these features may be reduced under cattle grazing. Herbivores affect soil by compacting it where they tread and altering its nutrient status where they produce dung and urine. This directly affects soil insects and arachnids and indirectly affects foliar insects and arachnids by changing plant species composition through changes in soil status, as opposed to forage selection during grazing. Trampling is seen as generally harmful to the arthropod fauna ⁶⁴, although dung deposition provides a niche for additional species²⁸.

The patterns of indicative beetle species from the geostatistical analysis³², and management and environmental correlations of their distributions using direct gradient analysis, allowed a classification of the broader assemblage of 73 species of ground and rove beetles (Fig. 6). A total of 28% of the beetle assemblage was sensitive to the effects of grazing (indicated by partitioned slice), 40% to landform, 28% sensitive to an interaction between grazing and landform and with the remaining 4% unallocated (Fig. 6). The smaller satellite circles indicate the direction of the response of these beetle species to each factor (Fig. 6). Sanderson et al.⁴⁶ provide further evidence of the importance that abiotic factors have in affecting the spatial patterns of arthropod populations in upland landscapes at larger spatial scales. Soil moisture or site wetness is recognized as a major determining factor in the distribution of many ground beetle species¹⁰.

Conclusions

There were significant effects of grazing on the species of various arthropod taxa, consistent with other investigations ^{47,48}. Sensitivity to grazing management is related to taxon and trophic level. The most marked direct effects were on small, herbivorous leafhoppers. Vegetation structure was important for small, herbivorous species and these indirect effects from grazing also influenced the distribution of webs and species composition of webbuilding spiders. The effects of grazing on the larger, ground-active beetles, namely ground and rove beetles, and wolf spiders was not consistent. About a third of beetle species significantly responded to differences in grazing management. However, landform was a considerable influence on these wider ranging, generalist predators. Overall, it is possible to distinguish between those species primarily sensitive to land use (represented by grazing) and to landform (particularly as it relates to climate change). Clearly, botanical composition and structural heterogeneity are important considerations in the objective to conserve or enhance upland biodiversity, but grazing is not the sole driver. Landscapescale studies are necessary to place into context detailed information on mammalian herbivore - vegetation - arthropod interactions.

The diversity of ground beetles, plant hoppers (Homoptera: Auchenorhyncha) and spiders (Araneae) in grassland, heathland and montane ecosystems is related to botanical diversity and the structural variability of vegetation ^{13, 49,50,46 30,58}. For grasslands in general, there is a positive correlation between the number of botanical species and the species richness of bees (Hymenoptera: Apidoidea), butterflies (Lepidoptera), phytophagous beetles (Coleoptera: Chrysomelidae) and true bugs (Hemiptera)⁴⁸. Differences between insect and arachnid species in the effects of grazing are highlighted by work undertaken in Wytham Woods, Oxfordshire ^{51,42,47}. For spider assemblages, changes in plant architecture were found to be most important, while other species (e.g. leaf miners) were affected more by variations in floristic species composition. As a consequence, leaf miners had a rapid turnover whereas spider species accumulated over time. Age of grassland was also important; there are common species of spiders, leaf miners and leaf hoppers (Cicadellidae) that are restricted to old grasslands.

The structural complexity of tussock-forming grasses encourages more planthoppers and web-building spiders in upland grasslands ⁵². For wolf spiders (Araneae: Lycosidae), species which pursue prey and do not build webs, the reverse is true and ca. 92% of individuals of this family mainly use the shorter grass between tussocks ⁵⁴. The relationships between grazers, vegetation and arthropods for upland, indigenous grasslands were consistent with the situation in lowland grasslands. The diversity of many arthropod taxa of lowland grasslands were favoured primarily by an increase in average vegetation height ^{53; 42;47, 55}.

Although general arthropod diversity is diminished by intensive management, many species respond within a few years to favourable changes in vegetation structure and botanical species composition that result from modified management ^{56,51,42,57,30}. Similarly, Coleoptera (beetles) of lowland grasslands are affected by the intensity of pasture management and species of ground beetles (Carabidae) have been identified that are resilient to or excluded by such management ^{36,57,10}. The indicator species of intensively managed, lowland pastures are typically smaller ³³ and more mobile ⁵⁸ than those of less intensively managed pastures. However, the inheritance effects of drainage, fertilizer and lime inputs, and the cultivated pasture may constrain the benefits to arthropod diversity of altered management of intensified pastures compared with indigenous grasslands.

To conserve arthropod diversity in grasslands used for livestock production, whether lowland or upland, rotational grazing management is advocated ^{12,59,60,29,32}. This would provide habitat for those species requiring short swards and intense grazing, as well as large, diverse invertebrate assemblages in the taller, older swards. A mosaic including patches of grassland that have been ungrazed for two to 15 years may be optimum for conservation of grassland invertebrates ^{61,12}. This is comparable with the conser-

vation management of Dutch salt marsh grasslands recommended by van Wieren⁶², that grazing should be between the level at which all grass is grazed short, and the level at which parts of the grassland become rank. In such conditions the grassland naturally becomes a mosaic of areas of short and tall grass and it was recommended for the Dutch grasslands that management should achieve an optimum ratio of 1:1 between these components. Further research is required to assess whether such mosaics are appropriate to sustain populations of various arthropod species typical of upland grasslands and heaths.

Acknowledgement

Iain Gordon, Mark Young and Richard Aspinall also worked on this research, funded by the Scottish Executive Environment and Rural Affairs Department and its predecessors, 1993-2003. Additional assistance in fieldwork and sample analysis was provided by Chris Bentley, Paola Sabbatini, Damien Cagnard, Gordon Common, Jim Small and Alison Smith. Martin Luff, Peter Hammond and Mike Davidson kindly carried out verification of the species identifications. Thanks to Pat Carnegie for assistance with the illustrations and to Robin Pakeman for useful comments on the first draft.

References

- ¹Fuller, R.J. and Gough, S.J. 1999. Changes in sheep numbers in Britain: implications for bird populations. Biological Conservation **91**: 73-89.
- ²Miles, J. 1988. Vegetation and soil changes in the uplands. In: Usher, M.B. and Thompson, D.B.A. (eds.). Ecological Change in the Uplands. Oxford: Blackwell Scientific Publications. p. 57-70.
- ³Grant, S.A. and Armstrong, H.M. 1993. Grazing ecology and the conservation of heather moorland: the development of models as aids to management. Biodiversity and Conservation 2: 79-94.
- ⁴Clarke, J. L., Welch, D. and Gordon, I. J. 1995. The influence of vegetation pattern on the grazing of heather moorland by red deer and sheep. I. The location of animals on grass/heather mosaics. Journal of Applied Ecology **32**: 166-176.
- ⁵Hurlbert, I.A.R., Iason, G.R. and Racey, P.A. 1996. Habitat utilisation in a stratified upland landscape by two lagomorphs with different feeding strategies. Journal of Applied Ecology **33**: 315-324.
- ⁶Milne, J.A., Birch, C.P.D., Hester, A.J., Robertson, A., Aspinall, R.J., Dennis, P. and Armstrong, H.M. 1996. The impact of vertebrate herbivores on the natural heritage of the Scottish Uplands - a review (Chapter 5. Impact on invertebrates). Report to Scottish Natural Heritage and the Scottish Office Agriculture, Environment and Fisheries Department, Edinburgh. p. 127.
- ⁷Tudor, G.J. and Mackey, E.C. 1995. Upland land cover change in post-war Scotland. In: Thompson, D.B.A., Hester, A.J. and Usher, M.B. (eds.). Heaths and Moorland: Cultural Landscapes. Edinburgh: HMSO. p. 28-42.
- ⁸Ratcliffe, D.A. (Ed.) 1977. A Nature Conservation Review. Cambridge: Cambridge University Press. p. 417.
- ⁹McLean, I.F.G. 1990. The fauna of calcareous grasslands. In: Hillier, S.H., Walter, D.W.H. and Wells, D.A. (eds.). Calcareous grasslands ecology and management. Huntingdon, UK: Bluntisham Books. p. 41-46.
- ¹⁰Rushton, S.P., Luff, M.L. and Eyre, M.D. 1991. Habitat characteristics of grassland *Pterostichus* species (Coleoptera: Carabidae). Ecological Entomology **16**: 91-104.
- ¹¹Rodwell, J.S. (Ed.) 1992. British Plant Communities. Volume 3. Grasslands and montane communities. Cambridge: Cambridge University Press. p. 540.
- ¹²Morris, M.G. 1991. The management of reserves and protected areas. In: Spellerberg, F.F., Goldsmith, F.B. and Morris, M.G. (eds.). The scien-

tific management of temperate communities for conservation. Oxford: Blackwell Publications. p. 323-347.

- ¹³ Coulson, J. C. and Whittaker, J. B. 1978. Ecology of moorland animals. In: Heal, O.W., Perkins, D.F. (eds.). Production Ecology of British Moors and Montane Grasslands. Berlin: Springer-Verlag. p. 52-93.
- ¹⁴ Bauer, L.J. 1989. Moorland beetle communities on limestone habitat islands. I. Isolation, invasion and local species diversity in carabids and staphylinids. Journal of Animal Ecology 58: 1077-1098.
- ¹⁵Watkinson, A.R. and Ormerod, S.J. 2001. Grasslands, grazing and biodiversity: editors' introduction. Journal of Applied Ecology **38**: 233-237.
- ¹⁶Anon. 1995. Biodiversity: the UK Steering Group report. Meeting the Rio challenge and action plans. London: Her Majesty's Stationery Office. 103 p.
- ¹⁷Perlman, D.L. and Adelson, G. 1997. Biodiversity. Exploring Values and Priorities in Conservation. New York: J. Wiley and Sons. p. 182.
- ¹⁸ Lawton, J.H. 1994. What do species do in ecosystems? Oikos **71**: 367-374.
- ¹⁹ Ribera, I., Doledec, S., Downie, I.S. and Foster, G.N. 2001. Effect of land disturbance and stress on species traits of ground beetle assemblages. Ecology 82: 1112-1129.
- ²⁰Galbraith, H., Murray, S., Duncan, K., Smith, R., Whitfield, D.P. and Thompson, D.B.A. 1993. Diet and habitat use of the Dotterel *Charadrius morinellus* in Scotland Ibis **135**: 148-155.
- ²¹Park, K.J., Robertson, P.A., Campbell, S.T., Foster, R., Russel, Z.M., Newborn, D. and Hudson, P.J. 2001. The role of invertebrates in the diet, growth and survival of red grouse (*Lagopus lagopus scoticus*) chicks. Journal of Zoology **254**: 137-145.
- ²²Rebecca, G.W. and Bainbridge, I.P. 1998. The breeding status of the Merlin *Falco columbarius* in Britain in 1993-94. Bird Study **45**: 172-187.
- ²⁴Hester, A.J., Gordon, I.J., Baillie, G.J. and Tappin, E. 1999. Foraging behaviour of sheep and red deer within natural heather-grass mosaics Journal of Applied Ecology 36: 133-146.
- ²⁵Wallis de Vries, M.F. and Dalebondt, C. 1994. Foraging strategy of cattle in patchy grassland. Oecologia **100**: 98-106.
- ²⁶Oom, S.P., Hester, A.J., Elston, D.A. and Legg, C.J. 2002. Spatial interaction models: from human geography to plant-herbivore interactions. Oikos 98: 65-74.
- ²⁷Coulson, J.C. and Butterfield, J.E.L. 1985. The invertebrate communites of peat and upland grasslands in the north of England and some conservation implications. Biological Conservation **34**: 197-225.
- ²⁸Coulson, J.C. 1988. The structure and importance of invertebrate communities on peatlands and moorlands, and effects of environmental changes. In: Usher M.B. and Thompson D.B.A. (eds.). Ecological change in the uplands. Oxford: Blackwell Scientific Publications. p. 365-380.
- ²⁹Dennis, P., Young, M.R., Howard, C.L. and Gordon, I.J. 1997. The response of epigeal beetles (Col.: Carabidae; Staphylinidae) to varied grazing regimes on upland *Nardus stricta* grasslands. Journal of Applied Ecology **34**: 433-443.
- ³⁰Dennis, P., Young, M.R. and Gordon, I.J. 1998. Distribution and abundance of small insects and arachnids in relation to structural heterogeneity of grazed, indigenous grasslands. Ecological Entomology 22: 253-264.
- ³¹Dennis, P., Young, M.R. and Bentley, C. 2001. The effects of varied grazing management on epigeal spiders, harvestmen and pseudoscorpions of *Nardus stricta* grassland in upland Scotland. Agriculture, Environment and Ecosystems 86: 39-57.
- ³²Dennis, P., Aspinall, R.J. and Gordon, I.J. 2002. Spatial distribution of upland beetles in relation to landform, vegetation and grazing management. Basic and Applied Ecology 3: 183-193.
- ³³Blake, S., Foster, G.N., Eyre, M.D. and Luff, M.L. 1994. Effects of habitat type and grassland management practices on the body size distribution of carabid beetles. Pedobiologia 38: 502-512.
- ³⁴ Rushton, S.P., Luff, M.L. and Eyre, M.D. 1989. Effects of pasture improvement and management on the ground beetle and spider communities of upland grasslands. Journal of Applied Ecology 26: 489-503.
- ³⁵Gardner, S.M., Hartley, S.E., Davies, A. and Palmer, S.C.F. 1997. Carabid

communities on heather moorlands in north-east Scotland: the consequences of grazing pressure for community diversity. Biological Conservation **81**: 275-286.

- ³⁶Eyre, M., Luff, M., Rushton, S. and Topping, C. 1989. Ground beetles and weevils (Carabidae and Curculionidae) as indicators of grassland management practices. Journal of Applied Entomology **107**: 508-517.
- ³⁷McFerran, D.M., Montgomery, W.I. and McAdam, J.H. 1994a. Effects of grazing intensity on heathland vegetation and ground beetle assemblages of the uplands of County Antrim, north-east Ireland. Biology and Environment: Proceedings of the Royal Irish Academy **94B**: 41-52.
- ³⁸McFerran, D.M., Montgomery, W.I. and McAdam, J.H. 1994b. The impact of grazing on communities of ground-dwelling spiders (Araneae) in upland vegetation types. Biology and Environment: Proceedings of the Royal Irish Academy **94B**: 119-126.
- ³⁹Perring, F. 1959. Topographical gradients of chalk grassland. Journal of Ecology 47: 447-481.
- ⁴⁰Whittaker, J.B. and Tribe, N.P. 1998. Predicting numbers of an insect (*Neophilaenus lineatus*: Homoptera) in a changing climate. Journal of Animal Ecology **67**: 987-991.
- ⁴¹Waloff, N. 1980. Studies on grassland leafhoppers (Auchenorrhyncha: Homoptera) and their natural enemies. Advances in Ecological Research 11: 82-215.
- ⁴²Gibson, C.W.D., Brown, V.K., Losito, L. and Gavin, GC. 1992a. The response of invertebrate assemblies to grazing. Ecography 15: 166-176.
- ⁴³Hulme, P.D., Pakeman, R.J., Torvell, L., Fisher, J.M. and Gordon, I.J. 1999. The effects of controlled sheep grazing on the dynamics of upland *Agrostis-Festuca* grassland. Journal of Applied Ecology**36**: 886-900.
- ⁴⁴Grant, S.A., Torvell, L., Sim, E.M., Small, J.L. and Armstrong, R.H. 1996a. Controlled grazing studies on *Nardus* grassland. I. Effects of betweentussock sward height and species of grazer on *Nardus* utilisation and floristic composition in two fields in Scotland. Journal of Applied Ecology **33**: 1053-1064.
- ⁴⁵Grant, S.A., Torvell, L., Sim, E.M., Small, J.L. and Elston, D.A. 1996b. Seasonal patterns of leaf growth and senescence of *Nardus stricta* (L.) and responses of tussocks to differing severity, timing and frequency of defoliation. Journal of Applied Ecology **33**: 1145-1155.
- ⁴⁶ Sanderson, R.A., Rushton, S.P., Cherrill, A.J. and Byrne, J.P. 1995. Soil, vegetation and space: an analysis of their effects on the invertebrate communities of a moorland in north-east England. Journal of Applied Ecology **32**: 506-518.
- ⁴⁷Gibson, C.W.D., Hambler, C. and Brown, V.K. 1992b. Changes in spider (Araneae) assemblages in relation to succession and grazing management. Journal of Applied Ecology **29**: 132-142.
- ⁴⁸Tscharntke, T. and Greiler, H.-J. 1995. Insect communities, grasses and grasslands. Annual Review of Entomology **40**: 535-558.
- ⁴⁹Cherrill, A.J. and Rushton, S.P. 1993. The Auchenorhyncha of an unimproved moorland in northern England. Ecological Entomology 18: 95-103.
- ⁵⁰Downie, I.S., Butterfield, J.E.L. and Coulson, J.C. 1995. Habitat preferences of sub-montane spiders in northern England. Ecography 18: 51-61.
- ⁵¹Brown, V.K., Gibson, C.W.D. and Sterling, P.H. 1990. The mechanisms controlling insect diversity in calcareous grasslands. In: Hillier, S.H., Walton D.W.H. and Wells D.A. (eds.). Calcareous grasslands - ecology and management. Bluntisham, UK: Bluntisham Books. p. 79-87.
- ⁵²Cherrett, J.M. 1964. The distribution of spiders on the Moor House National Nature Reserve, Westmorland. Journal of Animal Ecology **33**: 27-48.
- ⁵³Morris, M.G. 1971. Differences between the invertebrate faunas of grazed and ungrazed chalk grasslands. IV. Abundance and diversity of Homoptera: Auchenorhyncha. Journal of Applied Ecology 8: 37-52.

⁵⁴Bayram, A. and Luff, M.L. 1993. Winter abundance and diversity of lycosids

(Lycosidae: Araneae) and other spiders in grass tussocks in a field margin. Pedobiologia **37**: 357-364.

- ⁵⁵Kruess, A. and Tscharntke, T. 2002. Contrasting responses of plant and insect diversity to variation in grazing intensity. Biological Conservation **106**: 293-302.
- ⁵⁶Duffey, E., Morris, M.G., Sheail, J., Ward, L.K., Wells, D.A. and Wells, T.C.E. 1974. Grassland Ecology and Wildlife Management. London: Chapman and Hall. p. 215.
- ⁵⁷Curry, J.P. 1994. Grasslands invertebrates. Ecology, influence on soil fertility and effects on plant growth. London: Chapman and Hall. p. 424.
- ⁵⁸Siemann, E., Tilman, D. and Haarstad, J. 1999. Abundance, diversity and body size: patterns from a grassland arthropod community Journal of Animal Ecology 68: 824-835.
- ⁵⁹Morris, M.G. and Rispin, W.E. 1988. A beetle fauna of oolitic limestone grassland, and the responses of species to conservation management by different cutting regimes. Biological Conservation 43: 87-105.
- ⁶⁰Morris, M.G. 1990. The effects of management on the invertebrate community of calcareous grassland. In: Hillier, S.H., Walton, D.W.H. and Wells, D.A. (eds.). Calcareous grasslands - ecology and management. Harpenden, UK: Bluntisham Books. p. 128-133.
- ⁶¹Thomas, J.A. 1990. The conservation of Adonis blue and Lulworth skipper butterflies - two sides of the same coin. In: Hillier, S.H., Walter, D.W.H. and Wells, D.A. (eds.). Calcareous Grasslands: Ecology and Management. Huntingdon, UK: Bluntisham Books. p. 112-117.
- ⁶²Van Wieren, S.E. 1991. The management of populations of large mammals. In: Spellerberg, F.F., Goldsmith, F.B. and Morris, M.G. (eds.). The scientific management of temperate communities for conservation. Oxford: Blackwell Scientific Publications. p. 103-128.
- ⁶³Getis, A. and Ord, J.K. 1992. The analysis of spatial association by use of distance statistics. Geographical Analysis 24: 189-206.
- ⁶⁴Usher, M.B. and Gardner, S.M. 1988. Animal communities in the uplands: how is naturalness influenced by management? In: Usher, M.B. and Thompson, D.B.A. (eds.). Ecological change in the uplands. Oxford: Blackwell Scientific Publications. p. 75-92.